*Neuroscience Letters*, 133 (1991) 315–318
1991 Elsevier Scientific Publishers Ireland Ltd.



## AUTHOR INDEX

Ang, L.C., see Juurlink, B.H.J., 25
Arvin, B., see Moncada, C., 287

Balaguer, L., Romano, J. and Ruiz-Pesini, P., Serotonin immunoreactivity in the autonomic intrapulmonary ganglia of the fetal sheep, 151
Baldissera, F., Cavallari, P. and Dworzak, F., Cramps: a sign of motoneurone 'bistability' in a human patient, 303
Ballinger, M.L., see Sheller, R.A., 113
Barajas, L., Liu, L. and Nishiyama, C., Prenatal and postnatal development of the CGRP-immunoreactive innervation in the rat kidney, 219
Bar-Peled, O., Israeli, M., Ben-Hur, H., Hoskins, I., Groner, Y. and Biegon, A., Developmental pattern of muscarinic receptors in normal and Down's syndrome fetal brain — an autoradiographic study, 154
Bateson, A.N., see Glencorse, T.A., 45
Bean, A.J., Xu, Z., Chai, S.Y., Brimijoin, S. and Hökfelt, T., Effect of intracerebral injection of monoclonal acetylcholinesterase antibodies on cholinergic nerve terminals in the rat central nervous system, 145
Beck, S.G. and Choi, K.C., 5-Hydroxytryptamine hyperpolarizes CA3 hippocampal pyramidal cells through an increase in potassium conductance, 93
Behan, M., Kroker, A. and Bolz, J., Cortical barrelfields in organotypic slice cultures from rat somatosensory cortex, 191
Ben-Hur, H., see Bar-Peled, O., 154
Berge, O.-G., see Mjellem-Joly, N., 121
Biegon, A., see Bar-Peled, O., 154
Bing, G., see Stone, E.A., 33
Bittner, G.D., see Sheller, R.A., 113
Boksa, P., see Chaudieu, I., 267
Bolz, J., see Behan, M., 191
Borrione, P., Cervella, P., Geuna, S., Giacobini-Robecchi, M.G., Poncino, A. and Silengo, L., Electrophoretic analysis of neuronal genomic DNA from hypertrophic spinal ganglia during lizard tail regeneration, 245
Bouras, C., Vallet, P.G. and Hof, P.R., Asymmetric increase in substance P immunoreactivity in the rat and guinea pig substantia nigra after unilateral neocortical ablation, 53
Bourque, C.W., see Kirkpatrick, K., 271
Brimijoin, S., see Bean, A.J., 145
Bringmann, A. and Klingberg, F., Depth distribution of frontal cortical field potential to acoustic stimulation in unrestrained non-anaesthetized rats, 179
Brown, D.A., see Marsh, S.J., 298
Bruel, A., Cherqui, G., Columelli, S., Margelin, D., Roudier, M., Sinet, P.-M., Prieur, M., Pérignon, J.-L. and Delabar, J., Reduced protein kinase C activity in sporadic Alzheimer's disease fibroblasts, 89
Bures, J., see Gallo, M., 187

Campbell, K.J., see Takada, M., 137
Capsoni, S., see Stankov, B., 68
Cassell, M.D. and Roberts, L., Ultrastructural evidence for an olfactory-autonomic pathway through the rat central amygdaloid nucleus, 100
Cavallari, P., see Baldissera, F., 303
Cervella, P., see Borrione, P., 245
Chabot, J.-G., see Kar, S., 117
Chai, S.Y., see Bean, A.J., 145
Chan, J.Y.H., Lee, H.-C. and Chan, S.H.H., Reduction in postsynaptic $\alpha_2$-adrenoceptor activity by endogenous angiotensin III in the nucleus reticularis gigantocellularis of the rat, 81
Chan, S.H.H., see Chan, J.Y.H., 81
Chapouthier, G., see Martin, B., 73
Chartier-Harlin, M.-C., see Crawford, F., 1
Chaudieu, I., Mount, H., Quirion, R. and Boksa, P., Transient postnatal increases in excitatory amino acid binding sites in rat ventral mesencephalon, 267
Cherqui, G., see Bruel, A., 89
Chiba, H., Mitamura, T., Fujisawa, S., Ogata, A., Aimoto, Y., Tashiro, K. and Kobayashi, K., Apolipoproteins in rat cerebrospinal fluid: a comparison with plasma lipoprotein metabolism and effect of aging, 207
Choi, K.C., see Beck, S.G., 93
Columelli, S., see Bruel, A., 89
Cooper, R.M., see McIntosh, A.R., 311
Corda, M.G., Orlandi, M., Deiana, A.M., Salvadori, S. and Giorgi, O., Age-related decrease in the density of benzodiazepine recognition sites in the brain of the fresh water eel (*Anguilla anguilla*), 168
Cos, S., see Crespo, D., 253
Cowburn, R.F., see O'Neill, C., 15
Cozzi, B., see Stankov, B., 68
Crawford, F., Hardy, J., Mullan, M., Goate, A., Hughes, D., Fidani, L., Roques, P., Rossor, M. and Chartier-Harlin, M.-C., Sequencing of exons 16 and 17 of the $\beta$-amyloid precursor protein gene in 14 families with early onset Alzheimer's disease fails to reveal mutations in the $\beta$-amyloid sequence, 1
Cremer, J.E., see Myers, R., 20
Crespo, D., Cos, S., Fernandez-Viadero, C. and Gonzalez, C., Ultrastructural changes in hypothalamic supraoptic nucleus neurons of ovariectomized estrogen-deprived young rats, 253
Crisp, T., see Monroe, P.J., 229
Crouch, M.F., see Hendry, I.A., 29
Cuénod, M., see Piani, D., 159

Darlison, M.G., see Glencorse, T.A., 45
Deiana, A.M., see Corda, M.G., 168
Del Arco, C., see Pascual, J., 279
Del Olmo, E., see Pascual, J., 279
Delabar, J., see Bruel, A., 89
DeLaPaz, R., see Steinberg, G.K., 225
Desforges, C., see Martin, B., 73
Díaz, A., see Pascual, J., 279
Do, K.Q., see Piani, D., 159
Dworzak, F., see Baldissera, F., 303

Fauteck, J., see Stankov, B., 68
Fernandez-Viadero, C., see Crespo, D., 253
Fidani, L., see Crawford, F., 1
Fite, K.V., see Montgomery, N.M., 105
Florack, V.J., see Watt, C.B., 86
Fodor, M. and Palkovits, M., Neuropeptide Y-containing neuronal

pathway from the spinal trigeminal nucleus to the pontine peribrachial region in the rat, 195
Folstein, S.E., see Hedreen, J.C., 257
Fontana, A., see Piani, D., 159
Fowler, C.J., see O'Neill, C., 15
Frackowiak, R.S.J., see Myers, R., 20
Fraschini, F., see Stankov, B., 68
Fredow, G., see Richter, A., 57
Frei, K., see Piani, D., 159
Fujii, T., see Komori, K., 203
Fujisawa, S., see Chiba, H., 207
Fumagalli, P., see Stankov, B., 68
Furukawa, S., see Nakajima, M., 49

Gallo, M. and Bures, J., Acquisition of conditioned taste aversion in rats is mediated by ipsilateral interaction of cortical and mesencephalic mechanisms, 187
Geisert Jr., E.E., Murphy, T.P., Irwin, M.H. and Larjava, H., A novel cell adhesion molecule, G-CAM, found on cultured rat glia, 262
Geuna, S., see Borrione, P., 245
Ghez, C., see Martin, J.H., 61
Giacobini-Robecchi, M.G., see Borrione, P., 245
Gilbert, K.R., Aizenman, E. and Reynolds, I.J., Oxidized glutathione modulates *N*-methyl-D-aspartate- and depolarization-induced increases in intracellular $Ca^2$ in cultured rat forebrain neurons, 11
Giorgi, O., see Corda, M.G., 168
Glencorse, T.A., Bateson, A.N., Hunt, S.P. and Darlison, M.G., Distribution of the $GABA_A$ receptor α1- and γ2-subunit mRNAs in chick brain, 45
Goate, A., see Crawford, F., 1
Gollapudi, L., see Hentall, I.D., 215
González, A.M., see Pascual, J., 279
Gonzalez, C., see Crespo, D., 253
Gonzalez-Lima, F., see McIntosh, A.R., 311
Grafe, P., see Strupp, M., 237
Grant, G.A., see Steinberg, G.K., 225
Groner, Y., see Bar-Peled, O., 154
Gupta, B.B.P., Spessert, R. and Vollrath, L., Acetylcholine and muscarinic agonists increase synaptic ribbon numbers in the rat pineal, 125

Hardy, J., see Crawford, F., 1
Hattori, T., see Takada, M., 137
Hayashi, Y., see Nakajima, M., 49
Hedreen, J.C., Peyser, C.E., Folstein, S.E. and Ross, C.A., Neuronal loss in layers V and VI of cerebral cortex in Huntington's disease, 257
Hendry, I.A. and Crouch, M.F., Retrograde axonal transport of the GTP-binding protein $G_{i_2}$: a potential neurotrophic intra-axonal messenger, 29
Hentall, I.D., Kim, J.L. and Gollapudi, L., Responses of neurons in the ventromedial midbrain to noxious mechanical stimuli, 215
Hirano, A., see Kato, T., 163
Hof, P.R., see Bouras, C., 53
Hökfelt, T., see Bean, A.J., 145
Hole, K., see Mjellem-Joly, N., 121
Honoré, T., see Judge, M.E., 291
Hope, J., see Scott, J.R., 141
Hoskins, I., see Bar-Peled, O., 154
Hotta, H., Nishijo, K., Sato, A., Sato, Y. and Tanzawa, S., Stimulation of lumbar sympathetic trunk produces vasoconstriction of the vasa nervorum in the sciatic nerve via α-adrenergic receptors in rats, 249
Hughes, D., see Crawford, F., 1
Hunt, S.P., see Glencorse, T.A., 45

Irwin, M.H., see Geisert Jr., E.E., 262
Israeli, M., see Bar-Peled, O., 154
Iwasa, K.H., Li, M., Jia, M. and Kachar, B., Stretch sensitivity of the lateral wall of the auditory outer hair cell from the guinea pig, 171

Jaarsma, D., Sebens, J.B. and Korf, J., Localization of NMDA and AMPA receptors in rat barrel field, 233
Jacobsen, P., see Judge, M.E., 291
Jia, M., see Iwasa, K.H., 171
John, S.M., see Stone, E.A., 33
Judge, M.E., Sheardown, M.J., Jacobsen, P. and Honoré, T., Protection against post-ischemic behavioral pathology by the α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) antagonist 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo(f)quinoxaline (NBQX) in the gerbil, 291
Juurlink, B.H.J., Munoz, D.G. and Ang, L.C., Motoneuron survival in vitro: effects of pyruvate, α-ketoglutarate, gangliosides and potassium, 25

Kachar, B., see Iwasa, K.H., 171
Kar, S., Chabot, J.-G. and Quirion, R., Quantitative autoradiographic localisation of [$^{125}$I]endothelin-1 binding sites in spinal cord and dorsal root ganglia of the rat, 117
Kashiwagi, K., see Nakajima, M., 49
Katagiri, T., see Kato, T., 163
Kato, T., Hirano, A., Manaka, H., Sasaki, H., Katagiri, T., Kawanami, T., Shikama, Y., Seino, T. and Sasaki, H., Calcitonin gene-related peptide immunoreactivity in familial amyotrophic lateral sclerosis, 163
Kawanami, T., see Kato, T., 163
Kawanishi, K., see Niimi, M., 183
Kawashima, T., see Nakajima, M., 49
Kawatani, M., see Matsumoto, G., 211
Khalansky, A.S., see Preobrazhensky, A.A., 284
Kim, J.L., see Hentall, I.D., 215
Kingery, W.S., see Vallin, J.A., 241
Kirkpatrick, K. and Bourque, C.W., Dual role for calcium in the control of spike duration in rat supraoptic neuroendocrine cells, 271
Klingberg, F., see Bringmann, A., 179
Kobayashi, K., see Chiba, H., 207
Kobayashi, K., see Nakajima, M., 49
Kogure, K., see Nagasawa, H., 129
Komori, K., Fujii, T. and Nagatsu, I., Do some tyrosine hydroxylase-immunoreactive neurons in the human ventrolateral arcuate nucleus and globus pallidus produce only L-DOPA?, 203
Korf, J., see Jaarsma, D., 233
Kozlova, M.V., see Preobrazhensky, A.A., 284
Kroker, A., see Behan, M., 191
Kunis, D.M., see Steinberg, G.K., 225
Kupfermann, I., see Teyke, T., 307
Kurata, K. and Shibata, R., Effects of $D_1$ and $D_2$ antagonists on the transient increase of dopamine release by dopamine agonists by means of brain dialysis, 77

Larjava, H., Geisert Jr., E.E., 262
Lee, H.-C., see Chan, J.Y.H., 81
Li, M., see Iwasa, K.H., 171
Li, Z., see Montgomery, N.M., 105
Liu, L., see Barajas, L., 219
Lo, E.H., see Steinberg, G.K., 225
Löscher, W., see Richter, A., 57

Lucini, V., see Stankov, B., 68
Lund, A., see Mjellem-Joly, N., 121
Lux, H.D., see Mironov, S.L., 175

Manaka, H., see Kato, T., 163
Manjil, L.G., see Myers, R., 20
Margelin, D., see Bruel, A., 89
Marsh, S.J. and Brown, D.A., Potassium currents contributing to action potential repolarization in dissociated cultured rat superior cervical sympathetic neurones, 298
Martin, B., Desforges, C. and Chapouthier, G., Comparisons between patterns of convulsions induced by two $\beta$-carbolines in 10 inbred strains of mice, 73
Martin, J.H. and Ghez, C., Impairments in reaching during reversible inactivation of the distal forelimb representation of the motor cortex in the cat, 61
Mata, N.L., see Sun, J.-H., 97
Matsumoto, G., Kawatani, M. and Takeshige, C., Inhibitory effect of 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX) on the micturition reflex in rat, 211
McIntosh, A.R., Cooper, R.M. and Gonzalez-Lima, F., Metabolic activation of the rat visual system by patterned light and footshock, 311
Meldrum, B.S., see Moncada, C., 287
Mienville, J.-M., Comparison of fast responses to serotonin and 2-methyl-serotonin in voltage-clamped N1E-115 neuroblastoma cells, 41
Minors, D.S., Waterhouse, J.M. and Wirz-Justice, A., A human phase–response curve to light, 36
Mironov, S.L. and Lux, H.D., Calmodulin antagonists and protein phosphatase inhibitor okadaic acid fasten the 'run-up' of high-voltage activated calcium current in rat hippocampal neurones, 175
Mitamura, T., see Chiba, H., 207
Mitsumoto, Y., see Ohkubo, T., 275
Mjellem-Joly, N., Lund, A., Berge, O.-G. and Hole, K., Potentiation of a behavioural response in mice by spinal coadministration of substance P and excitatory amino acid agonists, 121
Mohri, T., see Ohkubo, T., 275
Moncada, C., Arvin, B., Peillet, E.L. and Meldrum, B.S., Non-NMDA antagonists protect against kainate more than AMPA toxicity in the rat hippocampus, 287
Monroe, P.J., Perschke, S.E., Crisp, T. and Smith, D.J., Evaluation of the interactions of serotonergic and adrenergic drugs with $\mu$, $\delta$, and $\kappa$ opioid binding sites, 229
Montgomery, N.M., Fite, K.V. and Li, Z., Anatomical evidence for an intergeniculate leaflet in *Rana pipiens*, 105
Mount, H., see Chaudieu, I., 267
Mullan, M., see Crawford, F., 1
Munoz, D.G., see Juurlink, B.H.J., 25
Murphy, T.P., see Geisert Jr., E.E., 262
Myers, R., Manjil, L.G., Frackowiak, R.S.J. and Cremer, J.E., [$^3$H]PK 11195 and the localisation of secondary thalamic lesions following focal ischaemia in rat motor cortex, 20

Nagasawa, H. and Kogure, K., Alterations of [$^3$H]inositol 1,4,5-trisphosphate binding in the postischemic rat brain, 129
Nagatsu, I., see Komori, K., 203
Nakajima, M., Kashiwagi, K., Hayashi, Y., Saito, M., Kawashima, T., Furukawa, S. and Kobayashi, K., $\alpha$-Tocopherol supports the survival and neurite extension of neurons cultured from various regions of fetal rat brain, 49
Niimi, M., Takahara, J., Sato, M. and Kawanishi, K., Neurotensin and growth hormone-releasing factor-containing neurons projecting to the median eminence of the rat: a combined retrograde tracing and immunohistochemical study, 183
Nishiyama, C., see Barajas, L., 219
Nishijo, K., see Hotta, H., 249

Ogata, A., see Chiba, H., 207
Ohkubo, T., Mitsumoto, Y. and Mohri, T., Characterization of the uptake of adenosine by cultured rat hippocampal cells and inhibition of the uptake by xanthine derivatives, 275
Onaka, T. and Yagi, K., Sino-aortic baroreceptors are not involved in the potentiation of vasopressin or oxytocin secretion observed after noxious stimuli in the rat, 133
O'Neill, C., Cowburn, R.F., Wiehager, B., Alafuzoff, I., Winblad, B. and Fowler, C.J., Preservation of 5-hydroxytryptamine$_{1A}$ receptor–G protein interactions in the cerebral cortex of patients with Alzheimer's disease, 15
Orlandi, M., see Corda, M.G., 168

Palkovits, M., see Fodor, M., 195
Pascual, J., Del Arco, C., González, A.M., Díaz, A., Del Olmo, E. and Pazos, A., Regionally specific age-dependent decline in $\alpha_2$-adrenoceptors: an autoradiographic study in human brain, 279
Pazos, A., see Pascual, J., 279
Peillet, E.L., see Moncada, C., 287
Pérignon, J.-L., see Bruel, A., 89
Perschke, S.E., see Monroe, P.J., 229
Peyser, C.E., see Hedreen, J.C., 257
Piani, D., Frei, K., Do, K.Q., Cuénod, M. and Fontana, A., Murine brain macrophages induce NMDA receptor mediated neurotoxicity in vitro by secreting glutamate, 159
Poljak, A., see Steinberg, G.K., 225
Poncino, A., see Borrione, P., 245
Popov, V.V. and Supin, A.Y., Interaural intensity and latency difference in the dolphin's auditory system, 295
Preobrazhensky, A.A., Khalansky, A.S. and Kozlova, M.V., Neurochordins of higher vertebrates: similarities in P-epitope-bearing polypeptide composition and a study of P-epitope expression in cultured neural tissue cells, 284
Prieur, M., see Bruel, A., 89
Puke, M.J.C., Xu, X.-J. and Wiesenfeld-Hallin, Z., Intrathecal administration of clonidine suppresses autotomy, a behavioral sign of chronic pain in rats after sciatic nerve section, 199

Quirion, R., see Chaudieu, I., 267
Quirion, R., see Kar, S., 117

Reekie, L.J.D., see Scott, J.R., 141
Reiter, R.J., see Sun, J.-H., 97
Reynolds, I.J., see Gilbert, K.R., 11
Richter, A., Fredow, G. and Löscher, W., Antidystonic effects of the NMDA receptor antagonists memantine, MK-801 and CGP 37849 in a mutant hamster model of paroxysmal dystonia, 57
Roberts, L., see Cassell, M.D., 100
Romano, J., see Balaguer, L., 151
Roques, P., see Crawford, F., 1
Ross, C.A., see Hedreen, J.C., 257
Rossor, M., see Crawford, F., 1
Roudier, M., see Bruel, A., 89
Rozental, R., In vitro denervation of frog skeletal muscle: expression of several conductance classes of nicotinic receptors, 65
Ruiz-Pesini, P., see Balaguer, L., 151

Saito, M., see Nakajima, M., 49
Salvadori, S., see Corda, M.G., 168
Sasaki, H., see Kato, T., 163
Sasaki, S., see Watanobe, H., 7
Sato, A., see Hotta, H., 249
Sato, M., see Niimi, M., 183
Sato, Y., see Hotta, H., 249
Scott, J.R., Reekie, L.J.D. and Hope, J., Evidence for intrinsic control of scrapie pathogenesis in the murine visual system, 141
Sebens, J.B., see Jaarsma, D., 233
Seino, T., see Kato, T., 163
Sheardown, M.J., see Judge, M.E., 291
Sheller, R.A., Ballinger, M.L. and Bittner, G.D., Long-term survival of severed crayfish giant axons is not associated with an incorporation of glial nuclei into axoplasm, 113
Shibata, R., see Kurata, K., 77
Shikama, Y., see Kato, T., 163
Silengo, L., see Borrione, P., 245
Sinet, P.-M., see Bruel, A., 89
Smith, D.J., see Monroe, P.J., 229
Spessert, R., see Gupta, B.B.P., 125
Stankov, B., Cozzi, B., Lucini, V., Capsoni, S., Fauteck, J., Fumagalli, P. and Fraschini, F., Localization and characterization of melatonin binding sites in the brain of the rabbit (*Oryctolagus cuniculus*) by autoradiography and in vitro ligand–receptor binding, 68
Steinberg, G.K., Lo, E.H., Kunis, D.M., Grant, G.A., Poljak, A. and DeLaPaz, R., Dextromethorphan alters cerebral blood flow and protects against cerebral injury following focal ischemia, 225
Stephens, P.J., Differential effects of alcohols on the spike threshold of an identified motor axon in a crab (*Pachygrapsus crassipes*), 3
Stone, E.A., Zhang, Y., John, S.M. and Bing, G., c-*Fos* response to administration of catecholamines into brain by microdialysis, 33
Strupp, M. and Grafe, P., A chloride channel in rat and human axons, 237
Sun, J.-H., Reiter, R.J., Mata, N.L. and Tsin, A.T.C., Identification of 11-*cis*-retinal and demonstration of its light-induced isomerization in the chicken pineal gland, 97
Supin, A.Y., see Popov, V.V., 295

Takada, M., Campbell, K.J. and Hattori, T., Regional localization of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) uptake: mismatch between its uptake and neurotoxic sites, 137
Takahara, J., see Niimi, M., 183
Takebe, K., see Watanobe, H., 7
Takeshige, C., see Matsumoto, G., 211
Tanzawa, S., see Hotta, H., 249
Tashiro, K., see Chiba, H., 207
Taylor, C.P., see Vartanian, M.G., 109
Teyke, T., Weiss, K.R. and Kupfermann, I., Activity of identified cerebral neuron correlates with food-induced arousal in *Aplysia*, 307
Tsin, A.T.C., see Sun, J.-H., 97

Vallet, P.G., see Bouras, C., 53
Vallin, J.A. and Kingery, W.S., Adjacent neuropathic hyperalgesia in rats: a model for sympathetic independent pain, 241
Vartanian, M.G. and Taylor, C.P., Different stereoselectivity of the enantiomers of HA-966 (3-amino-1-hydroxy-2-pyrrolidinone) for neuroprotective and anticonvulsant actions in vivo, 109
Vollrath, L., see Gupta, B.B.P., 125

Watanobe, H., Sasaki, S. and Takebe, K., Evidence that intravenous administration of interleukin-1 stimulates corticotropin releasing hormone secretion in the median eminence of freely moving rats: estimation by push-pull perfusion, 7
Waterhouse, J.M., see Minors, D.S., 36
Watt, C.B. and Florack, V.J., A double-label study demonstrating that enkephalin and somatostatin are localized in separate populations of amacrine cells in the larval tiger salamander retina, 86
Weiss, K.R., see Teyke, T., 307
Wiehager, B., see O'Neill, C., 15
Wiesenfeld-Hallin, Z., see Puke, M.J.C., 199
Winblad, B., see O'Neill, C., 15
Wirz-Justice, A., see Minors, D.S., 36

Xu, X.-J., see Puke, M.J.C., 199
Xu, Z., see Bean, A.J., 145

Yagi, K., see Onaka, T., 133

Zhang, Y., see Stone, E.A., 33